

Scaling of Tropical-Cyclone Dissipation

Albert Ossó¹, Álvaro Corral² & Josep Enric Llebot¹

¹*Grup de Física Estadística,*

Facultat de Ciències,

Universitat Autònoma de Barcelona,

E-08193 Bellaterra, Barcelona, Spain

²*Centre de Recerca Matemàtica,*

Edifici Cc, Campus UAB,

E-08193 Bellaterra, Barcelona, Spain

(Dated: September 30, 2009)

The influence of climate variability and global warming on the occurrence of tropical cyclones (TC) is a controversial issue [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. Existing historical databases on the subject are not fully reliable [9, 11, 12, 13], but a more fundamental hindrance is the lack of basic understanding regarding the intrinsic nature of tropical cyclone genesis and evolution [14]. It is known that tropical cyclones involve more than a passive response to changing external forcing [15], but it is not clear which dynamic behaviour best describes them. Here we present a new approach based on the application of the power dissipation index (*PDI*), which constitutes an estimation of released energy [3], to individual tropical cyclones. A robust law emerges for the statistics of *PDI*, valid in four different ocean basins and over long time periods. In addition to suggesting a novel description of the physics of tropical cyclones in terms of critical phenomena [16, 17, 18], the law allows to quantify their response to changing climatic conditions, with an increase in the largest *PDI* values with sea surface temperature or the presence of El Niño phenomenon, depending on the basin under consideration. In this way, we demonstrate that the recent upswing in North Atlantic hurricane activity does not involve TCs quantitatively different from those in other sustained high-activity periods prior to 1970.

One important characterization of a complex phenomenon is given by the fluctuations in the “size” of the phenomenon over successive occurrences [16, 17, 18, 19]. We refer to neither spatial size (area, volume) nor something like the Saffir-Simpson category [20]; rather, we seek a physically relevant measure of released energy. For a tropical cyclone, a reasonable proxy for this energy has been proposed by Emanuel [3], using the *PDI*, defined as

$$PDI \equiv \sum_t v_t^3 \Delta t ,$$

where t denotes time and runs over the entire lifetime of the storm and v_t is the maximum sustained surface wind velocity at time t . In available best-track records, measurements are provided at intervals of $\Delta t = 6$ hours. Note that in this paper the *PDI* value is associated with an individual tropical cyclone, not with the total annual activity in some ocean basin [3].

We analyze tropical cyclone best-track records for several ocean basins: the North Atlantic and Northeastern Pacific (from the National Hurricane Center [21]), and the Northwestern Pacific and the Southern Hemisphere (from the Joint Typhoon Warning Center [22]). We exclude the North Indian Ocean, due to the small number of storms in the most reliable portion of its records. [For more details, see the Supplementary Information, including Fig. S2.]

We display in Fig. 1(a) the *PDI* probability density, $D(PDI)$, normalized in the usual way ($\int_0^\infty D(PDI)dPDI = 1$), for all four basins. The distributions include all storms occurring during an extended period, either 1966-2007 or 1986-2007 as indicated in the legend. All four distributions (given vertical offsets for clarity) can be characterized by a power-law decay in their central regions,

$$D(PDI) \propto 1/PDI^\alpha,$$

where the exponent α is in between 0.95 and 1.25 (Supplementary Information, including Table S1). Deviations from the power law at small *PDI* values can be attributed to the deliberate incompleteness of the records for “not significant” TCs, whereas the more rapid decay at large *PDI* values is associated with the finite size of the basin. That is, the storms with the largest *PDI* do not have enough room to last a longer time, as their tracks are limited by the size of the basin, which introduces a cutoff in the distribution (at its separation from the power-law fit, roughly) (Supplementary Information, Figs. S3-S7). Variations in the definition of the *PDI*, for example excluding times during which the storm attains subtropical or extratropical status, do not modify the shape of the *PDI* distribution; the results are also unchanged when restricted to storms that do not make continental landfall (Supplementary Information, Figs. S8 and S9).

The degree of similarity between the basins is truly remarkable, given the variety of formative processes at work. TCs in the Western Pacific (typhoons) develop principally from the monsoon, for example, while North Atlantic hurricanes are mainly associated with easterly waves (and the degree of association depends on the intensity of the hurricane) [8, 23]. In addition, each regional centre or agency follows different protocols in obtaining their data, using techniques which have gradually improved [9, 11, 12, 13].

Nevertheless, the shape of the *PDI* distribution is robust over long time periods. Figure 1(b) shows that this consistency holds over a period of at least 100 years in the North Atlantic (where the record is longest; corrections to the calibration of the maximum velocity do not alter this pattern, Supplementary Information, including Fig. S10). Thus, even though the database is certainly incomplete prior to the satellite era, and even more unreliable before aircraft reconnaissance began in 1944, the fraction of missed storms seems to be independent of *PDI*. This finding may appear counterintuitive until we consider the fact that a long-lasting tropical cyclone might be recorded as two shorter storms if its track is lost at some point. A power-law distribution is robust against such splitting of the data (Supplementary Information).

The existence of a simple statistical distribution that describes the whole spectrum of tropical

cyclone sizes in different basins over a long period of time (apart from incompleteness and finite size effects) reflects a startling degree of unity in the phenomenon; the small tropical depressions are described in the same way as the full developed most severe storms. Moreover, a robust power-law distribution is the hallmark of scale invariance [18]: there is no typical tropical cyclone *PDI*, up to the maximum allowed in a given ocean basin. The fact that all scales are equally important for energy dissipation poses a great challenge to the modelling of this complex phenomenon, and even to large-scale global climate simulations [15].

Scale invariance can occur in processes where perturbations propagate through a critical (i.e., highly susceptible) medium [16, 17, 19]. Thus, our result could indicate that the atmosphere, or perhaps the ocean-atmosphere system, is close to a critical state. In fact, this idea is not new; already in the 70's it was suggested that atmospheric convection takes place in a near-unstable environment [24]. Much more recently, Peters and Neelin have demonstrated the existence of a non-equilibrium stability-instability transition to which the state of the atmosphere is attracted [25]. Some properties of this transition can be obtained from static images of convecting cloud fields [26] or local observations of precipitation [27]. These findings support our complex-system approach to tropical-cyclone evolution; in correspondence, we provide a complementary perspective to the puzzle that these atmospheric processes constitute.

In addition, an important property of critical systems is that perturbations can evolve while keeping a delicate balance between growth and attenuation, resulting in sudden intensifications and deintensifications. Although recent years have seen considerable improvement in the prediction of tropical cyclone trajectories, reliable forecasts of their intensities have not yet been achieved [28, 29]. This failure may not be just due to technical limitations; it may be a fundamental feature of the criticality of tropical cyclone evolution.

Tropical cyclone activity shows large interannual variability. One important factor controlling such variability is sea surface temperature (*SST*). We average *SST* from the Hadley Center [30] over the same spatial areas and months in the TC season than Webster *et al.* [5] in order to get an annual (so, seasonal) *SST* value for each basin; then, we separate the *PDI* density into two contributions, one for years with *SST* above its long-term mean value $\langle SST \rangle$ (i.e., high *SST*) and another one for *SST* below $\langle SST \rangle$ (low *SST*). Mathematically, $\langle SST \rangle \equiv \sum_y SST(y)/Y$, where $SST(y)$ refers to year y and Y is the total number of years. In this way we eliminate the effect of interannual variations in the number of TCs and concentrate on a comparison of the individual tropical cyclones characterizing each type of year.

Remarkably, for the North Atlantic and the Northeastern Pacific, the resulting distributions have essentially the same shape as the distribution for all years but with a difference in scale: high-*SST* years are characterized by a larger value of the finite-size cutoff, and conversely for low-*SST* years, as can be seen in Fig. 2(a). The other two basins show much minor *PDI* variation with *SST*.

If we rescale each conditional distribution by a power of its mean value, $\langle PDI \rangle$, such that $PDI \rightarrow PDI / \langle PDI \rangle^\nu$ and $D(PDI) \rightarrow \langle PDI \rangle^\beta D(PDI)$, with $\nu = \beta = 1$ for $\alpha \leq 1$, and $\nu = 1/(2 - \alpha)$ and $\beta = \alpha/(2 - \alpha)$ for $\alpha > 1$, it becomes apparent that, for each basin, both distributions share a similar shape, as shown in Fig. 2(b). (The reason of this rescaling is the fact that a distribution with $\alpha > 1$ does not scale linearly with its mean value, see Supplementary Information.) Then, the difference between high-*SST* and low-*SST* *PDI* distributions rests mainly in the scale of the finite-size cutoff and not in the shape of the distribution.

Years with high *SST* are thus characterized by hurricanes with larger *PDI* values. As the *PDI* integrates the cube of the velocity over the storm lifetime, larger *PDI* values can result from longer lifetimes, larger (6-hour) velocities, or both. An analysis of the distributions of these variables shows that the increase in *PDI* comes mainly from an increase in the velocities, most apparent in the range above about 100 knots (i.e., corresponding to category 3 hurricanes and beyond [5]), in comparison with years of low *SST* (Supplementary Information, Fig. S11).

An analogous study can be done as a function of the so-called *MEI* index [31], which quantifies the strength of El Niño phenomenon. Taking annual values of *MEI*, years with $MEI > 0$ (corresponding to El Niño) lead to increased *PDI* values in the Northeastern and Northwestern Pacific, but keeping again the same shape of the *PDI* distribution, and the opposite for $MEI < 0$, see Figs. 2(a) and 2(b). In the case of the Northwestern Pacific, the contribution of TC durations to the increase in *PDI* is larger than in the rest of the cases (Supplementary Information). This is somehow related to the findings of Refs. [32, 33]. Very little variation with *MEI* is observed in the other two basins. Other indices (*AMO*, *NAO*, etc.) do not seem to influence the *PDI* distribution in any basin.

It is a well-known fact that individual years of high (or low) TC activity cluster into longer periods of predominantly high (or low) activity. For example, the North Atlantic has seen extraordinarily high activity between 1995 and 2005, which has been linked to global warming through an increase in the sea surface temperature [3, 5]. The issue is nonetheless controversial [4, 6, 11]. According to our analysis, the *PDI* distribution for the period 1995-2005 is indistinguishable from the distribution for years with high *SST* between 1966 and 2007, as well as from the distribution

for other periods of high activity, like 1926-1970 [1], see Fig. 3. We conclude that the recent dramatic increase of activity does not lead to unprecedented energy releases by individual TCs (although even higher SST could further increase the *PDI* values).

- [1] Goldenberg, S. B., Landsea, C. W., Mestas-Nuñez, A. M. & Gray, W. M. The Recent Increase in Atlantic Hurricane Activity: Causes and Implications. *Science* **293**, 474–479 (2001).
- [2] Trenberth, K. Uncertainty in Hurricanes and Global Warming. *Science* **308**, 1753–1754 (2005).
- [3] Emanuel, K. Increasing destructiveness of tropical cyclones over the past 30 years. *Nature* **436**, 686–688 (2005).
- [4] Landsea, C. W. Hurricanes and global warming. *Nature* **438**, E11–E12 (2005).
- [5] Webster, P. J., Holland, G. J., Curry, J. A. & Chang, H.-R. Changes in Tropical Cyclone Number, Duration, and Intensity in a Warming Environment. *Science* **309**, 1844–1846 (2005).
- [6] Chan, J. C. L. Comment on “Changes in Tropical Cyclone Number, Duration, and Intensity in a Warming Environment”. *Science* **311**, 1713b (2006).
- [7] Klotzbach, P. J. Trends in global tropical cyclone activity over the past twenty years (1986-2005). *Geophys. Res. Lett.* **33**, L10805 (2006).
- [8] Shepherd, J. M. & Knutson, T. The Current Debate on the Linkage Between Global Warming and Hurricanes. *Geography Compass* **1**, 1–24 (2007).
- [9] Kossin, J. P., Knapp, K. R., Vimont, D. J., Murnane, R. J. & Harper, B. A. A globally consistent reanalysis of hurricane variability and trends. *Geophys. Res. Lett.* **34**, L04815 (2007).
- [10] Elsner, J. B., Kossin, J. P. & Jagger, T. H. The increasing intensity of the strongest tropical cyclones. *Nature* **455**, 92–95 (2008).
- [11] Gray, W. M. Comments on “Increasing destructiveness of tropical cyclones over the past 30 years”. <http://arxiv.org> **0601050** (2006).
- [12] Landsea, C. W., Harper, B. A., Hoarau, K. & Knaff, J. A. Can We Detect Trends in Extreme Tropical Cyclones? *Science* **313**, 452–454 (2006).
- [13] Landsea, C. W. Counting Atlantic Tropical Cyclones Back to 1900. *Eos* **88** (**18**), 197–202 (2007).
- [14] Emanuel, K. *Divine wind: the history and science of hurricanes* (Oxford University Press, New York, 2005).
- [15] Emanuel, K. The Hurricane-Climate Connection. *Bull. Am. Met. Soc.* (**5**), ES10–ES20 (2008).

- [16] Bak, P. *How Nature Works: The Science of Self-Organized Criticality* (Copernicus, New York, 1996).
- [17] Turcotte, D. L. *Fractals and Chaos in Geology and Geophysics* (Cambridge University Press, Cambridge, 1997), 2nd edn.
- [18] Christensen, K. & Moloney, N. R. *Complexity and Criticality* (Imperial College Press, London, 2005).
- [19] Malamud, B. D. Tails of natural hazards. *Phys. World* **17** (8), 31–35 (2004).
- [20] Kantha, L. Time to Replace the Saffir-Simpson Hurricane Scale? *Eos* **87** (1), 3–6 (2006).
- [21] Jarvinen, B. R., Neumann, C. J. & David, M. A. S. A tropical cyclone data tape for the North Atlantic basin, 1886-1983: contents, limitations, and uses. <http://www.nhc.noaa.gov/pdf/NWS-NHC-1988-22.pdf> (1988).
- [22] Chu, J.-H., Sampson, C. R., Levine, A. S. & Fukada, E. The Joint Typhoon Warning Center Tropical Cyclone Best-Tracks, 1945-2000. https://metocph.nmci.navy.mil/jtwc/best_tracks/TC_bt_report.html (2002).
- [23] Landsea, C. W. A climatology of intense (or major) hurricanes. *Mon. Weather Rev.* **121**, 1703–1713 (1993).
- [24] Arakawa, A. & Schubert, W. H. Interaction of a cumulus cloud ensemble with the large-scale environment, part I. *J. Atmos. Sci.* **31**, 674–701 (1974).
- [25] Peters, O. & Neelin, J. D. Critical phenomena in atmospheric precipitation. *Nature Phys.* **2**, 393–396 (2006).
- [26] Peters, O., Neelin, J. D. & Nesbitt, S. W. Mesoscale convective systems and critical clusters. *J. Atmos. Sci.* (2009).
- [27] Peters, O., Hertlein, C. & Christensen, K. A Complexity View of Rainfall. *Phys. Rev. Lett.* **88**, 018701 (2002).
- [28] Willoughby, H. E. Forecasting Hurricane Intensity and Impacts. *Science* **315**, 1232–1233 (2007).
- [29] National Hurricane Center Forecast Verification. <http://www.nhc.noaa.gov/verification> (2008).
- [30] Rayner, N. A. *et al.* Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *J. Geophys. Res.* **108** (D14), 4407 (2003).
- [31] Wolter, K. & Timlin, M. S. Measuring the strength of ENSO events: how does 1997/98 rank? *Weather* **53**(9), 315–324 (1998).
- [32] Lander, M. A. An Exploratory Analysis of the relationship between Tropical Storm Formation in the Western North Pacific and ENSO. *Mon. Weather Rev.* **122**, 636–651 (1994).
- [33] Emanuel, K. Environmental Factors Affecting Tropical Cyclone Power Dissipation. *J. Clim.* **20**,

5497–5509 (2007).

Supplementary Information

Available.

Acknowledgements

We have benefited from the expertise and kindness of A. Deluca, K. Emanuel, E. Fukada, A. González, J. Kossin, B. Mathiesen, M. Paczuski, O. Peters and A. Turiel. A.O. and A.C. were put in contact through G. Orriols. The initial part of our research has been financed by the EXPLORA - INGENIO 2010 program, and also partially by grants by other grants of the Spanish MEC and Generalitat de Catalunya.

Author Information

Correspondence and requests for materials should be addressed to A.C. (ACorral at crm dot cat).


Competing financial interests

The authors declare no competing financial interests.


FIG. 1: | **Power-law distributions of tropical-cyclone PDI values.** (a) PDI probability densities for tropical cyclones in the North Atlantic, Northeastern Pacific, Northwestern Pacific, and Southern Hemisphere basins. The period considered is either 1966-2007 or 1986-2007, as indicated in the legend, depending on the reliability of the records and the sufficiency of statistics. The values in the vertical axis are divided by the factors 1, $\sqrt{1000}$, 1000 and $\sqrt{1000^3}$, to separate the curves for clarity. From top to bottom, the number of tropical cyclones in each case is 469, 674, 690 and 601. The distributions are consistent with a power law (straight lines) over most of their range, with exponents 1.19, 1.17, 0.98 and 1.11 (± 0.07 , Supplementary Information). The Kolmogorov-Smirnov (KS) test yields p -values larger than 60 % in all basins (Supplementary Information). Deviations from the power law at large PDI values reflect the finite size effect. (b) PDI probability densities for tropical cyclones in the North Atlantic over the 54-year periods 1900-1953 and 1954-2007, with 436 and 579 storms respectively. A two-sample KS test (Supplementary Information) gives a p -value around 15% (51% if $PDI > 2 \cdot 10^9 m^3/s^2$). The robustness of the distribution is apparent despite a notable lack of homogeneity in the quality of records.


FIG. 2: | **Scaling of PDI distributions conditioned to the values of SST and MEI.** (a) PDI probability densities calculated separately for years with high SST and for years with low SST, as well as for years with $MEI > 0$ and for years with $MEI < 0$. Tropical depressions (storms whose maximum v_t is below 34 knots, 1 knot = 1.85 km/h) are excluded from the Northwestern Pacific dataset, in order to give all basins the same treatment. Time periods and vertical offsets are as in Fig. 1(a). (b) As with the previous panel, except that the axes are rescaled by $\langle PDI \rangle^{\nu(\alpha)}$ and $1/\langle PDI \rangle^{\beta(\alpha)}$, with $\alpha = 1.19, 1.17, 1.17$ and 1.0 (the data set at the bottom has been shifted extra factors 100 and 10^4 in each axis for clarity sake); the two-sample KS test yields p -values equal to 90 %, 73 %, 44 % and 73 %, for $PDI/\langle PDI \rangle^\nu > 0.0003, 0.0005, 0.0005$ and 0.05 , in units of $(m^3/s^2)^{1-\nu}$; the ratios of mean PDI between high and low annual values of SST or MEI, $\langle PDI \rangle_{high}/\langle PDI \rangle_{low}$, are 1.4, 1.6, 1.4, and 1.4, always from top to bottom.

FIG. 3: | **Equivalence between periods of predominantly high activity and high SST.** PDI probability densities are compared for the years 1995-2005 and 1926-1970, as well as for years with high SST during 1966-2007. The number of storms is 166, 427 and 250, respectively. Two-sample KS tests yield p -values larger than 40% for each pair of distributions (Supplementary Information, Table S2); so we can conclude that no significant difference in the energy of TCs can be observed between these periods.



1(a)





2(a)

